

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



ALPHABETICAL
LIST

BIOACTIVE
PEPTIDES

IMMUNO-
CHEMICALS

MOLECULAR
BIOLOGY

RBI,
NEUROSCIENCE,
SIGNAL
TRANSDUCTION

TISSUE
CULTURE

OTHER
PRODUCT
GROUPS/USP

EQUIPMENT,
BOOKS AND
SUPPLIES

DIAGNOSTIC
KITS AND
REAGENTS

PRODUCT
INDEX

Biochemicals and Reagents

for Life Science Research

 **SIGMA®**

ALPHABETICAL LIST OF COMPOUNDS

PRODUCT NUMBER	DESCRIPTION	PRODUCT NUMBER	DESCRIPTION
D 1500	DIDANSYL-L-TYROSINE XOSE (C ₁₂ H ₁₄ N ₂ O ₄ S) FW: 310 mg	D 1500	(Continuation of D 1500) L-TYROSINE A
D 1500	Approx. 95% (HPLC)	D 1500	2',3'-DIDEOXYCYTIDINE 5'-TRIPHOSPHATE
CEC	Crystalline (xylene-C ₁₂ H ₁₄ N ₂ O ₄ S) FW: 612.8; fcc	CEC	0963-0
	[1263-03-2] C ₁₂ H ₁₄ N ₂ O ₄ S • FW: 612.8; fcc		Minimum 90% (HPLC)
D 2375	N,O-DIDANSYL-L-TYROSINE	D 2375	1 mg
CEC	Monocyclohexylammonium Salt	CEC	107.80
	Approx. 95%		FW: 451.2 (for free acid)
	[102783-47-1] C ₁₂ H ₁₄ N ₂ O ₄ S ₂ • C ₆ H ₁₃ N • FW: 746.9		R: 61-8: 45-36/37/39-22
D 6389	1,2-DIDÉCANOYL-β-D-GLYCEROL	D 6389	10 mg
C(10:0)	[(1,2-Dicapryl)glycerol] 10 mg	C(10:0)	32.00
CEC	Approx. 99%	CEC	116.35
	Cell-permeable activator of protein kinase C in platelets		Ref.: Lapetina, E.G., et al., J. Biol. Chem., 260 , 1358 (1985).
	[82950-64-9] C ₂₀ H ₄₀ O ₂ • FW: 400.6		R: 20/21/22 S: 36-22
D 1493	2',3-DIDEHYDRO-3'-DEOXY- THYMIDINE	D 1493	10 mg
CEC	(1',2,3-Dideoxy-β-D-glycero-pent-2-enylidene-2-enuronosyl)thymine; 2',3'-Anhydrotymidine; d4T)	CEC	109.65
	Minimum 98%		Ref.: Lin, T.S., et al., Biochem. Pharmacol., 17 , 2713 (1978).
	Nucleoside analog which inhibits HIV replication in vitro		Baba, M., et al., Biochem. Biophys. Res. Commun., 142 , 128 (1987).
	[3056-7-5] C ₁₀ H ₁₄ N ₂ O ₄ • FW: 224.2		[3056-7-5] C ₁₀ H ₁₄ N ₂ O ₄ • FW: 224.2
D 1285	2',3-DIDEOXYADENOSINE	D 1285	1 mg
DDA	[(DDA)-dA] 5 mg	DDA	71.10
CEC	Approx. 97%	CEC	225.60
	[4097-22-7] C ₁₀ H ₁₄ N ₂ O ₄ • FW: 235.2		Ref.: Sanger, F., et al., Proc. Natl. Acad. Sci. USA, 74 , 5463 (1977).
D 7284	2',3-DIDEOXYADENOSINE 5'-TRI- PHOSPHATE	D 7284	1 mg
CEC	Sodium Salt	CEC	69.50
	Minimum 85% (HPLC)		Inhibitor of DNA polymerase I - catalyzed chain elongation
	[See also: Molecular Biology Reagents Page 1610 [1326-96-5-9] C ₁₀ H ₁₄ N ₂ O ₄ P ₃] FW: 475.2 (for free acid)]		Ref.: Sanger, F., et al., Proc. Natl. Acad. Sci. USA, 74 , 5463 (1977).
	2',6-DIDEOXY-D-ALLOSE		[See also: Molecular Biology Reagents Page 1610 [1326-96-5-9] C ₁₀ H ₁₄ N ₂ O ₄ P ₃] FW: 475.2 (for free acid)]
D 6782	2',3-DIDEOXYCYTIDINE	D 6782	100 mg
DDC	[(DDC)-dC] 250 mg	DDC	72.80
CEC	Minimum 98%	CEC	160.20
	[7481-89-2] C ₁₀ H ₁₄ N ₂ O ₃ • FW: 211.2		500 mg
	FW: 211.2; See also: Immunochemicals Page 1304		288.70
	R: 61-8: 45-36/37/39-22		[102783-47-1] C ₁₀ H ₁₄ N ₂ O ₃ • FW: 211.2
D 7409	DIDEOXYCYTIDINE, ANTIBODY TO	D 7409	1 mg
	See: Immunochemicals Page 1304		112.80
D 7409	2',3-DIDEOXYCYTIDINE, ³ H-Labeled	D 7409	2 mg
	See: Immunochemicals Page 1305 and Radiochemicals Page 217		187.95
D 4770	2',3-DIDEOXYCYTIDINE 5'-TRIPHOSPHATE	D 4770	1 mg
CEC	Inhibitor of DNA polymerase I - catalyzed chain elongation	CEC	69.50
	Ref.: Sanger, F., et al., Proc. Natl. Acad. Sci. USA, 74 , 5463 (1977).		5 mg
	[See also: Molecular Biology Reagents Page 1610 [68726-28-3] C ₁₀ H ₁₄ N ₂ O ₄ P ₃] FW: 491.2 (for free acid)]		80.40
D 1542	Lithium Salt	D 1542	1 mg
CEC	Minimum 85%	CEC	144.75
	[93939-77-0] C ₉ H ₁₄ N ₂ O ₃ P ₃		(2-Hydroxymethyl)-3,4-pyrrolidinediol
	FW: 451.2 (for free acid)		α-Glucosidase inhibitor
	[R: 61-8: 45-36/37/39-22]		Ref.: Fleet, G.W., and Smith, P.W., Tetrahedron, 42 , 5685 (1986). [MATUSIK, R.J., et al., HCl GRW: 169.6 [100937-52-8] C ₉ H ₁₄ N ₂ O ₃ • HCl GRW: 169.6

(Continued)

To place an order call 800-325-3010 or visit www.sigma-aldrich.com/orders

MOLECULAR BIOLOGY PRODUCTS

Molecular Biology Products

PRODUCT NUMBER	DESCRIPTION	PRICE US \$	PRODUCT NUMBER
(Continuation of) PRIMER SETS; Fluorescent-Labeled for Automated Sequencing			
P 3348	(-40) M13 Forward Primer Set (5'-GTT-TTC CCA GTC ACG ACG-3')		
P 3223	(-29) M13 Reverse Primer Set (5'-CAG GAA ACA GCT ATG ACC-3')		
P 3473	Lambda gt10 Forward Primer Set (5'-AGC AAG TTC AGC CTG GTT AAG-3')		
P 6098	Lambda gt10 Reverse Primer Set (5'-CTT ATG AGT ATT TCT TCC AGG GTA-3')		
P 6723	T3 Phage Promoter Primer Set (5'-ATT AAC CCT CAC TAA AGG GA-3')		
P 7348	T7 Phage Promoter Primer Set (5'-TAA TAC GAC TCA CTA TAG GG-3')		
P 7973	SP6 Promoter Primer Set (5'-ATT-TAG GTG ACA CTA TAG-3')		
P 8598	Poly(T)AGC Primer Set (5'-TTT TTT TTT TTT TTT TTT TTA GC-3')		
NUCLEOTIDE SOLUTIONS FOR RESOLVING SEQUENCE AMBIGUITIES			
	Used for eliminating band compressions caused by the ability of G residues to form secondary structures which are not fully denatured during electrophoresis.		
	Ref.: 1. Mizusawa, S., et al., Nucl. Acids Res., 14, 1319 (1986).		
	2. Barr, P.J., et al., Biotechniques, 4, 428 (1986).		
D 5163	2'-Deoxyinosine 5'-Triphosphate (dITP) Sodium Salt 5 mM aqueous solution [95648-77-4] C ₁₀ H ₁₅ N ₄ O ₁₃ P ₃ FW 492.2 (for free acid)	1 μmole 15.50 Shipped in dry ice	
D 8783	7-Deaza-2'-deoxyguanosine 5'-Triphosphate (-N ⁷ -dGTP) Lithium Salt 10 mM aqueous solution [101515-08-6] C ₁₁ H ₁₇ N ₄ O ₁₃ P ₃ FW 506.2 (for free acid)	0.5 μmole 54.60 Shipped in dry ice	
DIDEOXYNUCLEOSIDE TRIPHOSPHATE SOLUTIONS 10 mM solutions; pH 7.0			
D 5413	2',3'-Dideoxyadenosine 5'-Triphosphate (ddATP) Lithium Salt [93939-70-9]	0.5 μmole 22.55 Shipped in dry ice	
D 5538	2',3'-Dideoxycytidine 5'-Triphosphate (ddCTP) Lithium Salt [93939-77-6] R: 61 S: 45-36/37/39-23	0.5 μmole 22.55 Shipped in dry ice	
D 5663	2',3'-Dideoxyguanosine 5'-Triphosphate (ddGTP) Lithium Salt [93939-69-6]	0.5 μmole 22.55 Shipped in dry ice	
D 1789	2',3'-Dideoxyinosine 5'-Triphosphate (ddITP) Lithium Salt [93858-64-1]	0.5 μmole 56.30 Shipped in dry ice	
D 5288	(3'-Deoxythymidine 5'-Triphosphate; ddTTP) (dTTP) Lithium Salt [93939-78-7]	0.5 μmole 22.55 Shipped in dry ice	
MANUAL AND AUTOMATED SEQUENCING KITS			
SEQ-1	DNA SEQUENCING REAGENTS FOR MAXAM-GILBERT (CHEMICAL DEGRADATION) METHODOLOGY This DNA sequencing kit contains all necessary buffers and modifying reagents used in the Maxam-Gilbert chemical degradation method of DNA sequencing. NO DEA LICENSE IS REQUIRED FOR THIS KIT. Kit contains sufficient reagents for approx. 100 μl of sequencing reactions.	1 kit	
D 5154	Dimethyl Sulfate Stop Solution Used to stop guanosine modification reaction		
D 5279	Dimethyl Sulfate Modification reagent for guanosine		
D 5404	Dimethyl Sulfate Buffer Buffer for guanosine modification reaction		
F 4011	Ferric Chloride, 3.0 M For disposal of hydrazine		
H 1764	Hydrazine Stop Solution Used to stop pyrimidine modification reaction		
H 2761	Hydrazine, Anhydrous Modification reagent for pyrimidines		
P 5881	Piperidine Cleaves DNA strands at modified bases		
F 4636	Formic Acid Modification reagent for purine bases		
S 8388	Sodium Acetate, 0.3 M Used in ethanol precipitation of DNA		
S 8513	Sodium Chloride, 5.0 M Inhibits modification of thymine by hydrazine		
S 8263	Sodium Hydroxide, 5.0 M For disposal of dimethyl sulfate		
W 4502	Water, 18 megohm, 0.2 μm filtered For reconstitution of buffers and use in modifications reactions		
Ref.: Maxam, A.M. and Gilbert, W., Meth. Enzymol., 65, 499 (1980). R: 11-45-46-26/27/28-34-42/43 S: 45-26-36/37/39-23			